

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

I declare this is my own work.

AS FURTHER MATHEMATICS

Paper 1

Time allowed: 1 hour 30 minutes

Materials

- You must have the AQA Formulae and statistical tables booklet for A-level Mathematics and A-level Further Mathematics.
- You should have a scientific calculator that meets the requirements of the specification. (You may use a graphical calculator.)

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you require extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do **not** write outside the box around each page or on blank pages.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice


- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
TOTAL	



Answer **all** questions in the spaces provided.

- 1** The complex number ω is shown below on the Argand diagram.



Which of the following complex numbers could be ω ?

Tick (✓) **one** box.

[1 mark]

$\cos(-2) + i \sin(-2)$

☐

$\cos(-1) + i \sin(-1)$

☐

$\cos(1) + i \sin(1)$

☐

$\cos(2) + i \sin(2)$

☐

- 2** Given that $f(x) = 3x - 1$ find the mean value of $f(x)$ over the interval $4 \leq x \leq 8$

Circle your answer.

[1 mark]

6

11

17

23



- 3** The matrix **M** represents a rotation about the x -axis.

$$\mathbf{M} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & a & \frac{\sqrt{3}}{2} \\ 0 & b & -\frac{1}{2} \end{bmatrix}$$

Which of the following pairs of values is correct?

Tick (✓) **one** box.

[1 mark]

$$a = \frac{1}{2} \quad \text{and} \quad b = \frac{\sqrt{3}}{2} \quad \boxed{}$$

$$a = \frac{1}{2} \quad \text{and} \quad b = -\frac{\sqrt{3}}{2} \quad \boxed{}$$

$$a = -\frac{1}{2} \quad \text{and} \quad b = \frac{\sqrt{3}}{2} \quad \boxed{}$$

$$a = -\frac{1}{2} \quad \text{and} \quad b = -\frac{\sqrt{3}}{2} \quad \boxed{}$$

- 4** The point $(2, -1)$ is invariant under the transformation represented by the matrix **N**

Which of the following matrices could be **N**?

Circle your answer.

[1 mark]

$$\begin{bmatrix} 4 & 6 \\ 2 & 5 \end{bmatrix}$$

$$\begin{bmatrix} 6 & 5 \\ 4 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 5 & 2 \\ 6 & 4 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 4 \\ 5 & 6 \end{bmatrix}$$

Turn over for the next question

Turn over ►



5

Show that the vectors $\begin{bmatrix} 1 \\ -3 \\ 5 \end{bmatrix}$ and $\begin{bmatrix} 7 \\ 4 \\ 1 \end{bmatrix}$ are perpendicular.

[2 marks]

6

Prove the identity

$$\cosh^2 x - \sinh^2 x = 1$$

[2 marks]



[3 marks]

[illegible]

Turn over ►



- 8** Stephen is correctly told that $(1 + i)$ and -1 are two roots of the polynomial equation

$$z^3 - 2iz^2 + pz + q = 0$$

where p and q are complex numbers.

- 8 (a)** Stephen states that $(1 - i)$ **must** also be a root of the equation because roots of polynomial equations occur in conjugate pairs.

Explain why Stephen's reasoning is wrong.

[1 mark]

- 8 (b)** Find p and q

[5 marks]



This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

Turn over ►



9 (a)

where a and b are constants to be determined.

[illegible]

9 (b)

[3 marks]

[illegible]

10 Matrix **A** is given by

$$\mathbf{A} = \begin{bmatrix} 3 & i-1 \\ i & 2 \end{bmatrix}$$

10 (a) Show that $\det \mathbf{A} = a + i$ where a is an integer to be determined.

[2 marks]

10 (b) Matrix **B** is given by

$$\mathbf{B} = \begin{bmatrix} 14-2i & b \\ c & d \end{bmatrix} \quad \text{and} \quad \mathbf{AB} = p\mathbf{I}$$

where $b, c, d \in \mathbb{C}$ and $p \in \mathbb{N}$

Find b, c, d and p

[6 marks]



[illegible]

Turn over ►



[1 mark]

$$\frac{1}{(r-1)!} - \frac{1}{r!} = \frac{r-1}{r!}$$

[3 marks]

$$\sum_{r=1}^n \frac{r-1}{r!} = a + \frac{b}{n!}$$

where a and b are integers to be determined.



[illegible]

Turn over ►



12 (a) Find the other two roots of the equation.

$$\cosh^3 \theta - 2 \cosh^2 \theta - \cosh \theta + 2 = 0$$

giving your answers in an exact form.

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



Prove by induction that, for all integers $n \geq 1$

$$\sum_{r=1}^n 2^{-r} = 1 - 2^{-n}$$

[illegible]

14 Curve C_1 has equation

$$\frac{x^2}{16} + \frac{y^2}{4} = 1$$

14 (a) Curve C_2 is a reflection of C_1 in the line $y = x$

Write down an equation of C_2

[1 mark]

14 (b) Curve C_3 is a circle of radius 4, centred at the origin.

Describe a single transformation which maps C_1 onto C_3


[2 marks]

14 (c) Curve C_4 is a translation of C_1
The positive x -axis and the positive y -axis are tangents to C_4

14 (c) (i) Sketch the graphs of C_1 and C_4 on the axes opposite. Indicate the coordinates of the x and y intercepts on your graphs.

[2 marks]





14 (c) (ii) Determine the translation vector.

[2 marks]

14 (c) (iii) The line $y = mx + c$ is a tangent to both C_1 and C_4
Find the value of m

[2 marks]

Turn over ►



$$\mathbf{r} = \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix} + \lambda \begin{bmatrix} 5 \\ 3 \\ -2 \end{bmatrix} \quad \text{and} \quad \frac{x-5}{4} = \frac{y}{2} = 4-z$$

[3 marks]

[illegible]


[1 mark]



[illegible]

16

Curve C has equation $y = \frac{ax}{x+b}$ where a and b are constants.
The equations of the asymptotes to C are $x = -2$ and $y = 3$

**16 (a)**

Write down the value of a and the value of b

[2 marks]

16 (b)

The gradient of C at the origin is $\frac{3}{2}$

With reference to the graph, explain why there is exactly one root of the equation

$$\frac{ax}{x+b} = \frac{3x}{2}$$

[2 marks]




$$\frac{ax}{x+b} \leq 1-x$$

[4 marks]

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

- 17** The curve C_1 has polar equation $r = 2a(1 + \sin \theta)$ for $-\pi < \theta \leq \pi$ where a is a positive constant.



The point M lies on C_1 and the initial line.

- 17 (a)** Write down, in terms of a , the polar coordinates of M

[1 mark]

- 17 (b)** N is the point on C_1 that is furthest from the pole O

Find, in terms of a , the polar coordinates of N

[2 marks]



[5 marks]

[illegible]


Turn over ►



17 (d) On the initial line below, sketch the graph of $r = 2a(1 + \cos \theta)$ for $-\pi < \theta \leq \pi$

Include the polar coordinates, in terms of a , of any intersection points with the initial line.

[2 marks]



END OF QUESTIONS



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



Do not write
outside the
box

[illegible]

[illegible]

[illegible]