

Centre No.						Paper Reference						Surname	Initial(s)	
Candidate No.						<b>6</b>	<b>6</b>	<b>6</b>	<b>6</b>	<b>/</b>	<b>0</b>	<b>1</b>	Signature	

Paper Reference(s)

**6666/01**

# Edexcel GCE

## Core Mathematics C4

### Advanced

Wednesday 26 January 2011 – Afternoon  
Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
<b>Total</b>	

**Materials required for examination**      **Items included with question papers**  
Mathematical Formulae (Pink)      Nil

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

**Instructions to Candidates**

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper. Answer ALL the questions. You must write your answer to each question in the space following the question. When a calculator is used, the answer should be given to an appropriate degree of accuracy.

**Information for Candidates**

A booklet 'Mathematical Formulae and Statistical Tables' is provided. Full marks may be obtained for answers to ALL questions. The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2). There are 7 questions in this question paper. The total mark for this paper is 75. There are 24 pages in this question paper. Any blank pages are indicated.

**Advice to Candidates**

You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner. Answers without working may not gain full credit.



*Turn over*



2. The current,  $I$  amps, in an electric circuit at time  $t$  seconds is given by

$$I = 16 - 16(0.5)^t, \quad t \geq 0$$

Use differentiation to find the value of  $\frac{dI}{dt}$  when  $t = 3$ .

Give your answer in the form  $\ln a$ , where  $a$  is a constant.

(5)







4. Relative to a fixed origin  $O$ , the point  $A$  has position vector  $\mathbf{i} - 3\mathbf{j} + 2\mathbf{k}$  and the point  $B$  has position vector  $-2\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ . The points  $A$  and  $B$  lie on a straight line  $l$ .

(a) Find  $\vec{AB}$ . (2)

(b) Find a vector equation of  $l$ . (2)

The point  $C$  has position vector  $2\mathbf{i} + p\mathbf{j} - 4\mathbf{k}$  with respect to  $O$ , where  $p$  is a constant. Given that  $AC$  is perpendicular to  $l$ , find

(c) the value of  $p$ , (4)

(d) the distance  $AC$ . (2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





5. (a) Use the binomial theorem to expand

$$(2-3x)^{-2}, \quad |x| < \frac{2}{3},$$

in ascending powers of  $x$ , up to and including the term in  $x^3$ . Give each coefficient as a simplified fraction.

(5)

$$f(x) = \frac{a+bx}{(2-3x)^2}, \quad |x| < \frac{2}{3}, \text{ where } a \text{ and } b \text{ are constants.}$$

In the binomial expansion of  $f(x)$ , in ascending powers of  $x$ , the coefficient of  $x$  is 0 and the coefficient of  $x^2$  is  $\frac{9}{16}$ . Find

(b) the value of  $a$  and the value of  $b$ ,

(5)

(c) the coefficient of  $x^3$ , giving your answer as a simplified fraction.

(3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---







Leave blank

6. The curve  $C$  has parametric equations

$$x = \ln t, \quad y = t^2 - 2, \quad t > 0$$

Find

(a) an equation of the normal to  $C$  at the point where  $t = 3$ , (6)

(b) a cartesian equation of  $C$ . (3)




Figure 1

The finite area  $R$ , shown in Figure 1, is bounded by  $C$ , the  $x$ -axis, the line  $x = \ln 2$  and the line  $x = \ln 4$ . The area  $R$  is rotated through  $360^\circ$  about the  $x$ -axis.

(c) Use calculus to find the exact volume of the solid generated. (6)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





7.

$$I = \int_2^5 \frac{1}{4 + \sqrt{x-1}} dx$$

- (a) Given that  $y = \frac{1}{4 + \sqrt{x-1}}$ , complete the table below with values of  $y$  corresponding to  $x = 3$  and  $x = 5$ . Give your values to 4 decimal places.

$x$	2	3	4	5
$y$	0.2		0.1745	

(2)

- (b) Use the trapezium rule, with all of the values of  $y$  in the completed table, to obtain an estimate of  $I$ , giving your answer to 3 decimal places.

(4)

- (c) Using the substitution  $x = (u - 4)^2 + 1$ , or otherwise, and integrating, find the exact value of  $I$ .

(8)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



