

2. A woman travels in a lift. The mass of the woman is 50 kg and the mass of the lift is 950 kg. The lift is being raised vertically by a vertical cable which is attached to the top of the lift. The lift is moving upwards and has constant deceleration of 2 m s^{-2} . By modelling the cable as being light and inextensible, find
- (a) the tension in the cable, **(3)**
- (b) the magnitude of the force exerted on the woman by the floor of the lift. **(3)**



3.




Figure 1

A box of mass 2 kg is held in equilibrium on a fixed rough inclined plane by a rope. The rope lies in a vertical plane containing a line of greatest slope of the inclined plane. The rope is inclined to the plane at an angle α , where $\tan \alpha = \frac{3}{4}$, and the plane is at an angle of 30° to the horizontal, as shown in Figure 1. The coefficient of friction between the box and the inclined plane is $\frac{1}{3}$ and the box is on the point of slipping up the plane. By modelling the box as a particle and the rope as a light inextensible string, find the tension in the rope.

(8)



Leave
blank

Question 3 continued

Lined area for writing the answer to Question 3.



Leave
blank

4. A lorry is moving along a straight horizontal road with constant acceleration. The lorry passes a point A with speed $u \text{ m s}^{-1}$, ($u < 34$), and 10 seconds later passes a point B with speed 34 m s^{-1} . Given that $AB = 240 \text{ m}$, find

(a) the value of u , (3)

(b) the time taken for the lorry to move from A to the mid-point of AB . (6)



5. A car is travelling along a straight horizontal road. The car takes 120 s to travel between two sets of traffic lights which are 2145 m apart. The car starts from rest at the first set of traffic lights and moves with constant acceleration for 30 s until its speed is 22 m s^{-1} . The car maintains this speed for T seconds. The car then moves with constant deceleration, coming to rest at the second set of traffic lights.

(a) Sketch, in the space below, a speed-time graph for the motion of the car between the two sets of traffic lights.

(2)

(b) Find the value of T .

(3)

A motorcycle leaves the first set of traffic lights 10 s after the car has left the first set of traffic lights. The motorcycle moves from rest with constant acceleration, $a \text{ m s}^{-2}$, and passes the car at the point A which is 990 m from the first set of traffic lights. When the motorcycle passes the car, the car is moving with speed 22 m s^{-1} .

(c) Find the time it takes for the motorcycle to move from the first set of traffic lights to the point A .

(4)

(d) Find the value of a .

(2)



6. A beam AB has length 15 m. The beam rests horizontally in equilibrium on two smooth supports at the points P and Q , where $AP = 2$ m and $QB = 3$ m. When a child of mass 50 kg stands on the beam at A , the beam remains in equilibrium and is on the point of tilting about P . When the same child of mass 50 kg stands on the beam at B , the beam remains in equilibrium and is on the point of tilting about Q . The child is modelled as a particle and the beam is modelled as a non-uniform rod.
- (a) (i) Find the mass of the beam.
- (ii) Find the distance of the centre of mass of the beam from A . **(8)**

When the child stands at the point X on the beam, it remains horizontal and in equilibrium. Given that the reactions at the two supports are equal in magnitude,

- (b) find AX . **(6)**



Question 6 continued

A series of 24 horizontal lines for writing.



P 4 1 8 2 8 A 0 1 9 2 8

7. [In this question, the horizontal unit vectors *i* and *j* are directed due east and due north respectively.]

The velocity, *v* m s⁻¹, of a particle *P* at time *t* seconds is given by

$$\mathbf{v} = (1 - 2t)\mathbf{i} + (3t - 3)\mathbf{j}$$

(a) Find the speed of *P* when *t* = 0 (3)

(b) Find the bearing on which *P* is moving when *t* = 2 (2)

(c) Find the value of *t* when *P* is moving

(i) parallel to *j*,

(ii) parallel to (-*i* - 3*j*). (6)



8.




Figure 2

Two particles A and B have masses $2m$ and $3m$ respectively. The particles are attached to the ends of a light inextensible string. Particle A is held at rest on a smooth horizontal table. The string passes over a small smooth pulley which is fixed at the edge of the table. Particle B hangs at rest vertically below the pulley with the string taut, as shown in Figure 2. Particle A is released from rest. Assuming that A has not reached the pulley, find

- (a) the acceleration of B , **(5)**
- (b) the tension in the string, **(1)**
- (c) the magnitude and direction of the force exerted on the pulley by the string. **(4)**



Leave blank

Question 8 continued

Lined writing area for the answer to Question 8.

Q8

(Total 10 marks)

TOTAL FOR PAPER: 75 MARKS

END

